


WHAT IS CLAIMED IS:

1. A compound of the Formula A:



5 wherein:

W=Y is selected from CR<sup>1</sup>=N, N=CR<sup>1</sup>,  $\text{C}(=\text{O})\text{NR}^{1'}$  or  $\text{R}^{1'}\text{N}-\text{C}(=\text{O})$ ;

10 a is 0 or 1; b is 0 or 1; m is 0, 1 or 2; p is 0, 1 or 2; q is 0, 1, 2 or 3; r is 0 or 1; s is 0 or 1; t is 2, 3, 4, 5 or 6;

Q is selected from: H, -NR<sup>6</sup>R<sup>7</sup>, aryl and heterocyclyl, said aryl and heterocyclyl which is optionally substituted with one to three R<sup>Z</sup>;

15 R<sup>1</sup> is independently selected from: 1) H, 2) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl, 3) (C=O)<sub>a</sub>O<sub>b</sub>aryl, 4) C<sub>2</sub>-C<sub>10</sub> alkenyl, 5) C<sub>2</sub>-C<sub>10</sub> alkynyl, 6) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl, 7) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl, 8) CO<sub>2</sub>H, 9) halo, 10) CN, 11) OH, 12) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, 13) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>6</sup>R<sup>7</sup>, 14) NR<sup>c</sup>(C=O)NR<sup>6</sup>R<sup>7</sup>, 15) S(O)<sub>m</sub>R<sup>a</sup>, 16) S(O)<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, 17) NR<sup>c</sup>S(O)<sub>m</sub>R<sup>a</sup>, 18) oxo, 19) CHO, 20) NO<sub>2</sub>, 21) NR<sup>c</sup>(C=O)O<sub>b</sub>R<sup>a</sup>, 22) O(C=O)O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl, 23) O(C=O)O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl, 24) O(C=O)O<sub>b</sub>aryl, 25) O(C=O)O<sub>b</sub>-  
20 heterocycle, and 26) O<sub>a</sub>-P=O(OH)<sub>2</sub>, said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R<sup>Z</sup>;

R<sup>1'</sup> is independently selected from: 1) H, 2) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl, 3) (C=O)<sub>a</sub>O<sub>b</sub>aryl, 4) C<sub>2</sub>-C<sub>10</sub> alkenyl, 5) C<sub>2</sub>-C<sub>10</sub> alkynyl, 6) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl, 7) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl, 8) CO<sub>2</sub>H, 9) halo, 10) CN, 11) OH, 12) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, 13) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>6</sup>R<sup>7</sup>, 14) S(O)<sub>m</sub>R<sup>a</sup>, 15) S(O)<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, 16) oxo, 17) CHO, 18) O(C=O)O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl, 19) O(C=O)O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl, 20)

O(C=O)O<sub>b</sub>aryl, 21) O(C=O)O<sub>b</sub>-heterocycle, and 22) O<sub>a</sub>-P=O(OH)<sub>2</sub>, said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R<sup>Z</sup>;

R<sup>2</sup> is independently selected from: 1) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl, 2) (C=O)<sub>a</sub>O<sub>b</sub>aryl, 3) C<sub>2</sub>-C<sub>10</sub> alkenyl, 4) C<sub>2</sub>-C<sub>10</sub> alkynyl, 5) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl, 6) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl, 7) CO<sub>2</sub>H, 8) halo, 9) CN, 10) OH, 11) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, 12) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>6</sup>R<sup>7</sup>, 13) NR<sup>c</sup>(C=O)NR<sup>6</sup>R<sup>7</sup>, 14) S(O)<sub>m</sub>R<sup>a</sup>, 15) S(O)<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, 16) NR<sup>c</sup>S(O)<sub>m</sub>R<sup>a</sup>, 17) CHO, 18) NO<sub>2</sub>, 19) NR<sup>c</sup>(C=O)O<sub>b</sub>R<sup>a</sup>, 20) O(C=O)O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl, 21) O(C=O)O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl, 22) O(C=O)O<sub>b</sub>aryl, 23) O(C=O)O<sub>b</sub>-heterocycle, and 24) O<sub>a</sub>-P=O(OH)<sub>2</sub>, said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one, two or three substituents selected from R<sup>Z</sup>;

R<sup>3</sup> and R<sup>4</sup> are independently selected from: H, C<sub>1</sub>-C<sub>6</sub>-alkyl and C<sub>1</sub>-C<sub>6</sub>-perfluoroalkyl, or

R<sup>3</sup> and R<sup>4</sup> are combined to form -(CH<sub>2</sub>)<sub>t</sub>- wherein one of the carbon atoms is optionally replaced by a moiety selected from O, S(O)<sub>m</sub>, -N(R<sup>b</sup>)C(O)-, and -N(COR<sup>a</sup>)-;

R<sup>5</sup> is independently selected from: 1) H, 2) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl, 3) (C=O)<sub>a</sub>O<sub>b</sub>aryl, 4) C<sub>2</sub>-C<sub>10</sub> alkenyl, 5) C<sub>2</sub>-C<sub>10</sub> alkynyl, 6) (C=O)<sub>a</sub>O<sub>b</sub> heterocyclyl, 7) (C=O)<sub>a</sub>O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl, 8) CO<sub>2</sub>H, 9) halo, 10) CN, 11) OH, 12) O<sub>b</sub>C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, 13) O<sub>a</sub>(C=O)<sub>b</sub>NR<sup>6</sup>R<sup>7</sup>, 14) NR<sup>c</sup>(C=O)NR<sup>6</sup>R<sup>7</sup>, 15) S(O)<sub>m</sub>R<sup>a</sup>, 16) S(O)<sub>2</sub>NR<sup>6</sup>R<sup>7</sup>, 17) NR<sup>c</sup>S(O)<sub>m</sub>R<sup>a</sup>, 18) oxo, 19) CHO, 20) NO<sub>2</sub>, 21) O(C=O)O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl, 22) O(C=O)O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and 23) O<sub>a</sub>-P=O(OH)<sub>2</sub>, said alkyl, aryl, alkenyl, alkynyl, heterocyclyl, and cycloalkyl optionally substituted with one or more substituents selected from R<sup>Z</sup>;

R<sup>6</sup> and R<sup>7</sup> are independently selected from: 1) H, 2) (C=O)O<sub>b</sub>R<sup>a</sup>, 3) C<sub>1</sub>-C<sub>10</sub> alkyl, 4) aryl, 5) C<sub>2</sub>-C<sub>10</sub> alkenyl, 6) C<sub>2</sub>-C<sub>10</sub> alkynyl, 7) heterocyclyl, 8) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, 9) SO<sub>2</sub>R<sup>a</sup>, 10) (C=O)NR<sup>b</sup>, 11) OH, and 12) O<sub>a</sub>-P=O(OH)<sub>2</sub>, said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one or more substituents selected from R<sup>Z</sup>, or

R<sup>6</sup> and R<sup>7</sup> can be taken together with the nitrogen to which they are attached to form a monocyclic or bicyclic heterocycle with 4-7 members in each ring and optionally containing, in addition to the nitrogen, one or more additional heteroatoms selected from N, O and S, said monocyclic or bicyclic heterocycle optionally substituted with one or more substituents selected from R<sup>Z</sup>;

R<sup>Z</sup> is selected from: 1) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>1</sub>-C<sub>10</sub>)alkyl, 2) O<sub>r</sub>(C<sub>1</sub>-C<sub>3</sub>)perfluoroalkyl, 3) (C<sub>0</sub>-C<sub>6</sub>)alkylene-S(O)<sub>m</sub>R<sup>a</sup>, 4) oxo, 5) OH, 6) halo, 7) CN, 8) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>2</sub>-C<sub>10</sub>)alkenyl, 9) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>2</sub>-C<sub>10</sub>)alkynyl, 10) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, 11) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-aryl, 12) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-heterocyclyl, 13) (C=O)<sub>r</sub>O<sub>s</sub>(C<sub>0</sub>-C<sub>6</sub>)alkylene-N(R<sup>b</sup>)<sub>2</sub>, 14) C(O)R<sup>a</sup>, 15) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>R<sup>a</sup>, 16) C(O)H, 17) (C<sub>0</sub>-C<sub>6</sub>)alkylene-CO<sub>2</sub>H, 18) C(O)N(R<sup>b</sup>)<sub>2</sub>, 19) S(O)<sub>m</sub>R<sup>a</sup>, 20) S(O)<sub>2</sub>N(R<sup>b</sup>)<sub>2</sub>, 21) NR<sup>c</sup>(C=O)O<sub>b</sub>R<sup>a</sup>, 22) O(C=O)O<sub>b</sub>C<sub>1</sub>-C<sub>10</sub> alkyl, 23) O(C=O)O<sub>b</sub>C<sub>3</sub>-C<sub>8</sub> cycloalkyl, 24) O(C=O)O<sub>b</sub>aryl, 25) O(C=O)O<sub>b</sub>-heterocycle, and 26) O<sub>a</sub>-P=O(OH)<sub>2</sub>, said alkyl, alkenyl, alkynyl, cycloalkyl, aryl, and heterocyclyl is optionally substituted with up to three substituents selected from R<sup>b</sup>, OH, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halogen, CO<sub>2</sub>H, CN, O(C=O)C<sub>1</sub>-C<sub>6</sub> alkyl, oxo, N(R<sup>b</sup>)<sub>2</sub> and O<sub>a</sub>-P=O(OH)<sub>2</sub>;


R<sup>a</sup> is: substituted or unsubstituted (C<sub>1</sub>-C<sub>6</sub>)alkyl, substituted or unsubstituted (C<sub>2</sub>-C<sub>6</sub>)alkenyl, substituted or unsubstituted (C<sub>2</sub>-C<sub>6</sub>)alkynyl, substituted or unsubstituted (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, substituted or unsubstituted aryl, (C<sub>1</sub>-C<sub>6</sub>)perfluoroalkyl, 2,2,2-trifluoroethyl, or substituted or unsubstituted heterocyclyl; and

R<sup>b</sup> is: H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, substituted or unsubstituted aryl, substituted or unsubstituted benzyl, substituted or unsubstituted heterocyclyl, (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, (C=O)OC<sub>1</sub>-C<sub>6</sub> alkyl, (C=O)C<sub>1</sub>-C<sub>6</sub> alkyl or S(O)<sub>2</sub>R<sup>a</sup>;

R<sup>c</sup> is selected from: 1) H, 2) C<sub>1</sub>-C<sub>10</sub> alkyl, 3) aryl, 4) C<sub>2</sub>-C<sub>10</sub> alkenyl, 5) C<sub>2</sub>-C<sub>10</sub> alkynyl, 6) heterocyclyl, 7) C<sub>3</sub>-C<sub>8</sub> cycloalkyl, and 8) C<sub>1</sub>-C<sub>6</sub> perfluoroalkyl, said alkyl, cycloalkyl, aryl, heterocyclyl, alkenyl, and alkynyl is optionally substituted with one or more substituents selected from R<sup>Z</sup>, or

or a pharmaceutically acceptable salt or a stereoisomer thereof.

2. The compound according to Claim 1 of the Formula B:



wherein:

$W=Y$  is selected from  $CR^1=N$ ,  $N=CR^1$ ;

5 or a pharmaceutically acceptable salt or a stereoisomer thereof.

3. The compound according to Claim 2 wherein:


10 Q is selected from:  $-NR^6R^7$ , phenyl and heterocyclyl which are optionally substituted with one to three  $R^Z$ ;

$R^a$  is:  $(C_1-C_6)$ alkyl,  $(C_3-C_6)$ cycloalkyl, aryl, or heterocyclyl; and

15  $R^b$  is: H,  $(C_1-C_6)$ alkyl, aryl, heterocyclyl,  $(C_3-C_6)$ cycloalkyl,  $(C=O)OC_1-C_6$  alkyl,  $(C=O)C_1-C_6$  alkyl or  $S(O)_2R^a$ ;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

4. The compound according to Claim 3 of the Formula C:



wherein:

Q is heterocyclyl, said heterocyclyl optionally substituted with 1 to 3  $R^Z$ ;

25  $R^2$  is independently selected from: 1)  $C_1-C_6$ alkyl, 2) aryl, 3) heterocyclyl, 4)  $CO_2H$ , 5) halo, 6) CN, 7) OH, 8)  $S(O)_2NR^6R^7$ , and 9)  $O_a-P=O(OH)_2$ , said alkyl, aryl and heterocyclyl optionally substituted with one, two or three substituents selected from  $R^Z$ ;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

## 5. A compound which is selected from:

- 1-(1-{4-[3-(1,3-oxazol-2-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 5 1-{1-[4-(6-phenyl-3-pyrimidin-2-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[3-(1H-imidazol-2-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 10 1-(1-{4-[3-(1-methyl-1H-pyrazol-5-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[6-phenyl-3-(1H-pyrazol-5-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[6-phenyl-3-(1H-pyrazol-5-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 15 1-(1-{4-[3-(1-methyl-1H-pyrazol-4-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[3-(1-methyl-1H-imidazol-2-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-[4-(6-phenyl-3-tetrahydrofuran-3-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 20 1-{1-[4-(6-phenyl-3-tetrahydrofuran-2-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-[4-(6-phenyl-3-thien-2-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 25 1-(1-{4-[3-(4-methylmorpholin-3-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[3-(1-acetylazetidin-3-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-[4-(6-phenyl-3-pyridin-2-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 30 1-{1-[4-(6-phenyl-3-pyridin-3-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-[4-(6-phenyl-3-pyridin-4-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;

- 1-(1-{4-[3-(morpholin-4-ylmethyl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[6-phenyl-3-(1,3-thiazol-2-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 5 1-(1-{4-[6-phenyl-3-(1,3-thiazol-5-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[6-phenyl-3-(1H-1,2,3-triazol-5-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 10 1-(1-{4-[6-phenyl-3-(1,3-thiazol-4-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[3-(1,1'-biphenyl-4-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-{1-[4-(2-methyl-3-oxo-6-phenyl-2,3-dihydro-1,2,4-triazin-5-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2H-benzimidazol-2-one;
- 15 1-{1-[4-(2-methyl-3-oxo-5-phenyl-2,3-dihydro-1,2,4-triazin-6-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2H-benzimidazol-2-one; and
- 1-(1-{4-[3-(methylthio)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 20 or a pharmaceutically acceptable salt or a stereoisomer thereof.

6. The TFA salt of a compound according to Claim 1 which is:

- 1-(1-{4-[3-(1,3-oxazol-2-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 25 1-{1-[4-(6-phenyl-3-pyrimidin-2-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl}-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[3-(1H-imidazol-2-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 30 1-(1-{4-[3-(1-methyl-1H-pyrazol-5-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[6-phenyl-3-(1H-pyrazol-5-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 35 1-(1-{4-[6-phenyl-3-(1H-pyrazol-5-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;

- 1-(1-{4-[3-(1-methyl-1H-pyrazol-4-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[3-(1-methyl-1H-imidazol-2-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 5 1-{1-[4-(6-phenyl-3-tetrahydrofuran-3-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl}-1,3-dihydro-2H-benzimidazol-2-one;
- 1-{1-[4-(6-phenyl-3-tetrahydrofuran-2-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl}-1,3-dihydro-2H-benzimidazol-2-one;
- 10 1-{1-[4-(6-phenyl-3-thien-2-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl}-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[3-(4-methylmorpholin-3-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[3-(1-acetylazetidin-3-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 15 1-{1-[4-(6-phenyl-3-pyridin-2-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl}-1,3-dihydro-2H-benzimidazol-2-one;
- 1-{1-[4-(6-phenyl-3-pyridin-3-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl}-1,3-dihydro-2H-benzimidazol-2-one;
- 20 1-{1-[4-(6-phenyl-3-pyridin-4-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl}-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[3-(morpholin-4-ylmethyl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[6-phenyl-3-(1,3-thiazol-2-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 25 1-(1-{4-[6-phenyl-3-(1,3-thiazol-5-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[6-phenyl-3-(1H-1,2,3-triazol-5-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 30 1-(1-{4-[6-phenyl-3-(1,3-thiazol-4-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-(1-{4-[3-(1,1'-biphenyl-4-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;
- 1-{1-[4-(2-methyl-3-oxo-6-phenyl-2,3-dihydro-1,2,4-triazin-5-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2H-benzimidazol-2-one;

1-{1-[4-(2-methyl-3-oxo-5-phenyl-2,3-dihydro-1,2,4-triazin-6-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2H-benzimidazol-2-one; and

1-(1-{4-[3-(methylthio)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;

or a stereoisomer thereof.

7. A compound according to Claim 5 which is selected from:

1-{1-[4-(6-phenyl-3-pyrimidin-2-yl-1,2,4-triazin-5-yl)benzyl]piperidin-4-yl}-1,3-dihydro-2H-benzimidazol-2-one;

1-(1-{4-[6-phenyl-3-(1H-1,2,3-triazol-5-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;

1-(1-{4-[6-phenyl-3-(1,3-thiazol-4-yl)-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one; and

1-(1-{4-[3-(1,1'-biphenyl-4-yl)-6-phenyl-1,2,4-triazin-5-yl]benzyl}piperidin-4-yl)-1,3-dihydro-2H-benzimidazol-2-one;

or a pharmaceutically acceptable salt or a stereoisomer thereof.

8. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 1.

9. A pharmaceutical composition comprising a pharmaceutical carrier, and dispersed therein, a therapeutically effective amount of a compound of Claim 5.

10. A method of inhibiting one or more of the isoforms of Akt in a mammal which comprises administering to the mammal a therapeutically effective amount of a compound of Claim 1.

11. A method of inhibiting one or more of the isoforms of Akt in a mammal which comprises administering to the mammal a therapeutically effective amount of a compound of Claim 5.

12. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 1.



13. A method for treating cancer which comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 5.

5 14. A method for treating a non-malignant disease in which angiogenesis is implicated which comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 1.

10 15. A method for treating a non-malignant disease in which angiogenesis is implicated which comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 5.

15 16. The composition of Claim 8 further comprising a second compound selected from: 1) an estrogen receptor modulator, 2) an androgen receptor modulator, 3) a retinoid receptor modulator, 4) a cytotoxic agent, 5) an antiproliferative agent, 6) a prenyl-protein transferase inhibitor, 7) an HMG-CoA reductase inhibitor, 8) an HIV protease inhibitor, 9) a reverse transcriptase inhibitor, 10) an angiogenesis inhibitor, 11) a PPAR- $\gamma$  agonist, 12) a PPAR- $\delta$  agonist, 13) an inhibitor of cell proliferation and survival signaling, and 14) an agent that interferes with a cell cycle checkpoint.

20 17. A method of treating cancer which comprises administering a therapeutically effective amount of a compound of Claim 1 in combination with radiation therapy.

25 18. A method of treating hyperproliferative disorders selected from restenosis, inflammation, autoimmune diseases and allergy/asthma which comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 1.

19. A method of treating hyperinsulinism which comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of Claim 1.